

State:Nagaland

Agriculture Contingency Plan for District: Kohima

1.0 District Agriculture profile				
1.1	Agro-Climatic/Ecological Zone			
	Agro Ecological Sub Region (ICAR)	Tropical to temperate		
	Agro-Climatic Zone (Planning Commission)	Eastern Himalayan Region		
	Agro Climatic Zone (NARP)	Sub – Tropical Hill Zone (98.10) & Mid Tropical Hill Zone (1.90) (The climate of this region is characterized by warm summer and mild winter with seasonal dry spells extending from November to April.)		
	List all the districts falling under the NARP Zone>(*>50% area falling in the zone)	Sub- tropical hill zone: Kohima, Mokokchung, Mon, Phek, Tuensang, Wokha, Zunheboto Mid-tropical Hill zone: Dimapur, Kohima, Mokokchung, Wokha		
	Geographic coordinates of district headquarters	Latitude	Longitude	Altitude
		25 ⁰ 40' N	94 ⁰ 08' E	1261 m (MSL)
	Name and address of the concerned ZRS/ ZARS/ RARS/ RRS/ RRTTS	ICAR, Nagaland centre, Jharnapani		
Mention the KVK located in the district with address	Krishi Vigyan Kendra, Kohima, Tesophenyu, 797109. Nagaland			

1.2	Rainfall	Normal RF (mm)	Normal Rainy days (number)	Normal Onset (specify week and month)	Normal Cessation (specify week and month)
	SW monsoon (3 rd week May - Sep):	1014.3	88	3 rd week May	4 th week September
	Post Monsoon/ NE Monsoon (Oct-Dec):	89.5	16	2 nd week October	4 th week December
	Winter (Jan- March)	61.2	12	1 st week January	4 th week March
	Summer (Apr-May)	470.2	29	2 nd week April	2 nd week May
	Annual	1653.2	145		

1.3	Land use pattern of the district (latest statistics)	Geographical area	Cultivable area	Forest area	Land under non-agricultural use	Permanent pastures	Cultivable wasteland	Land under Misc. tree crops and groves	Barren and uncultivable land	Current fallows	Other fallows
	Area ('000 ha)	146.300	58.420	72.194	4.390	0.500	6.900	9.350	5.190	21.426	14.706

* Source DAO office kohima 2017-18

1.4	Major Soils (common names like red sandy loam deep soils (etc.))*	Area ('000 ha)	Percent (%) of total
	1. Clay Soil	54.86	42.90%
	2. Loamy Soil	73.01	57.10%

* mention colour, depth and texture (heavy, light, sandy, loamy, clayey etc) and give vernacular name, if any, in brackets (data source: Soil Resource Maps of NBSS & LUP)

1.5	Agricultural land use	Area ('000 ha)	Cropping intensity %
	Net sown area	16.400	112.32
	Area sown more than once	2.020	
	Gross cropped area	18.42	

1.6	Irrigation	Area ('000 ha)		
	Net irrigated area	7.525		
	Gross irrigated area	7.580		
	Rainfed area	5.240		
	Sources of Irrigation	Number	Area ('000 ha)	Percentage of total irrigated area
	Canals (Community/Pvt Channel)	2571	6.085	
	Tanks			
	Open wells			
	Bore wells			
	Lift irrigation schemes			
	Micro-irrigation			
	Other sources (please specify)			

Total Irrigated Area			
Pump sets			
No. of Tractors			
Groundwater availability and use* (Data source: State/Central Ground water Department /Board)	No. of blocks/ Tehsils	(%) area	Quality of water (specify the problem such as high levels of arsenic, fluoride, saline etc)
Over exploited			
Critical			
Semi- critical			
Safe			
Wastewater availability and use			
Ground water quality	-		

*over-exploited: groundwater utilization > 100%; critical: 90-100%; semi-critical: 70-90%; safe: <70%

1.7 Area under major field crops & horticulture (statistical hand book of Nagaland 2017) (Specify year 2015-16)

1.7	S.No.	Major field crops cultivated	Area (*000 ha)							
			<i>Kharif</i>			<i>Rabi</i>			Summer	Grand total
			Irrigated	Rainfed	Total	Irrigated	Rainfed	Total		
1	Jhum Paddy	0.0	5.240	5.240	0.0	0.0	0.0	0.0	5.240	
2	TRC/WRC Paddy	0.0	10.110	10.110	-	-	-	-	10.110	
3	Maize	0.0	4.610	4.610	0.0	0.407	0.407	0.0	5.424	
4	Rapeseed & Mustard	0.0				0.203	0.203		0.203	
5	Sesamum	0.0	0.037	0.037					0.037	
6	Pea	0.0				.065	.065		.065	
7	Lentil	0.0								
8	Rajmah/kholar	0.0	.014	.014		.044	.044		.058	

S.No.	Horticulture crops - Fruits	Area ('000 ha)		
		Total	Irrigated	Rainfed
1	Banana	1.0		1.0
2	Passion fruit	1.1		1.1
3	Assam Lemon	0.1		0.1
4	Papaya	0.1		0.1
5	Plum	0.1		0.1
6	Orange	07		07
7	Pineapple	0.8		0.8
8	Colocasia	-		-
Others (specify)	Horticulture & plantation	4.58	0.0	4.58
	Horticulture crops - Vegetables	Total area ('000 ha)	Irrigated area ('000 ha)	Rainfed area ('000 ha)
1	Ginger	0.3	-	0.3
2	Cabbage	0.1	-	0.1
3	Potato	0.6	-	0.6
4	Tomato	0.2		0.2
Others (specify)	-			
	Medicinal and Aromatic crops	Total	Irrigated	Rainfed
1	Lemongrass	0.1	-	0.1
2	Citronella			
3	Neem			
4	Patchouli			
5	Amla			
Others (specify)	Spices	Total	Irrigated	Rainfed
1	Coriander			
2	Turmeric			

	3	Chilli			
	4	Ginger			
		Plantation crops	Total	Irrigated	Rainfed
	1	Coconut			
	2	Arecanut			
	Others (Specify)	Eg., industrial pulpwood crops etc.			
		Fodder crops	Total	Irrigated	Rainfed
	Others (Specify)	-			
		Total fodder crop area			
		Grazing land			
		Sericulture etc			
		Eri seeds (DFLS)			

1.8	Livestock (2012)	Male ('000)	Female ('000)	Total ('000)
	Non descriptive Cattle (local low yielding)	3261	7042	10.303
	Improved cattle	-	-	-
	Crossbred cattle	2975	8936	11.911
	Non descriptive Buffaloes (local low yielding)	0.935	0.912	1.847
	Pigs	-	-	53.928
	Goat	3.224	5.374	8.598
	Sheep	0.003	0.002	0.005
	Indigenous sheep	0.063	0.092	0.155
	Others (Camel, Pig, Yak etc.)	-	-	37.753
	Commercial dairy farms (Number)			
1.9	Poultry	No. of farms	Total No. of birds ('000)	
	Commercial (Poultry no)	-	198.408	
	Backyard			
	Duck	-	8.409	

1.10	Fisheries (Data source: Chief Planning Officer)						
	A. Capture						
	i) Marine (Data Source: Fisheries Department)	No. of fishermen	Boats		Nets		Storage facilities (Ice plants etc.)
			Mechanized	Non-mechanized	Mechanized (Trawl nets, Gill nets)	Non-mechanized (Shore Seines, Stake & trap nets)	
ii) Inland (Data Source: Fisheries Department)							
	B. Culture						
				Water Spread Area (ha)	Yield (t/ha)	Production ('000 tons)	
	i) Brackish water (Data Source: MPEDA/ Fisheries Department)						
	ii) Fresh water (Data Source: Fisheries Department)						
	Others						

1.11 Production and Productivity of major crops (2015-16)

1.11	Name of crop	Kharif		Rabi		Summer		Total		Crop residue as fodder ('000 tons)
		Production ('000 t)	Productivity (kg/ha)	Production ('000 t)	Productivity (kg/ha)	Production ('000 t)	Productivity (kg/ha)	Production ('000 t)	Productivity (kg/ha)	
Major Field crops (Crops to be identified based on total acreage)										
Crop 1	Jhum Paddy	10.33	1971	-	-	=	-	10.33	1971	
Crop 2	Rapeseed & Mustard			2.05	1009			2.05	1009	
Crop 3	Sesamum	0.24	648					0.24	628	
Crop 4	Pea			0.71	1092			0.71	1092	
Crop5	Lentil			0.15	833			0.15	833	
Crop6	Rajmah/khol ar			0.54	1227			0.54	1227	
Major Horticultural crops (Crops to be identified based on total acreage)										
Crop 1	Potato			16.18	10049			16.18	10049	
Crop 2	Rabi vegetables			15.661	18295			15.661	18295	
Crop 3	Kharif vegetables	20.165	9383					20.165	9383	
Crop 4	Arecanut									
Crop 5	Coconut									
Others	Banana	13.347	12024					13.347	12024	

1.12	Sowing window for 5 major field crops (start and end of normal sowing period)	Crop 1 : Jhum paddy	Crop 2: TRC/WRC Paddy	Crop 3: Maize	Crop 4: Turmeric & Ginger	Crop 5: Pea
	Kharif- Rainfed	April-May.	May-July	April-Aug.	April – May & July	-
	Kharif-Irrigated	-	-	-	-	-
	Rabi- Rainfed	-	-	-	-	Oct-Nov
	Rabi-Irrigated	-	-	-	-	-


1.13	What is the major contingency the district is prone to? (Tick mark)		Regular	Occasional	None	
	Drought		-		-	
	Flood		-	-		
	Cyclone		-	-		
	Hail storm		-		-	
	Heat wave		-	-		
	Cold wave			-	-	
	Frost		-	-		
	Sea water intrusion		-	-		
	Pests and disease outbreak (specify)		Rice-stem borer		-	-
			Rice blast disease	-		-
			Rhizome rot of ginger		-	-
Tomato (bacterial blight)				-	-	
Others (Landslides)			-		-	

1.14	Include Digital maps of the district for	Location map of district within State as Annexure I	Enclosed: Yes
		Mean annual rainfall as Annexure 2	Enclosed: Yes
		Soil map as Annexure 3	Enclosed: Yes

Annexure – 1: Location map of Kohima district



Annexure 3. Soil Map of Kohima district



2.0 Strategies for weather related contingencies

2.1 Drought –

2.1.1 Rainfed Situation

2.1.1.1 Pre- monsoon (2nd week of April to 2nd week of May).

Conditions			Suggested Contingency measures		
Early season drought (delayed onset)	Major Farming situation	Normal Crop / Cropping system	Change in crop / cropping system including variety	Agronomic measures	Remarks on Implementation
Delay by 2 weeks (2 nd to 3 rd week of April)	Moderately sloping on side slopes of hills with deep, fine loamy soils	Pre-kharif maize (local land races)	No change	No change	
	Gently sloping , side slopes of hills with moderately shallow fine soils	Jhum-Maize (local land races)	No change	No change	
	Steeply sloping hills with deep, fine soils	<i>Jhum</i> paddy (local land races)	No change	No change	
		Maize (local land races)	No change	No change	
	Moderately to gentle sloping hills slopes with deep loamy skeletal to fine loamy soils	<i>Jhum</i> paddy (local land races)	No change	No change	
		Jhum-Maize (local land races)	No change	No change	
Delay by 4 weeks (1 st to 2 nd week of May)	Moderately sloping on side slopes of hills with deep, fine loamy soils	Pre-kharif maize (local land races)	No change	No change	
	Gently sloping , side slopes of hills with moderately shallow fine soils	Jhum-Maize (local land races)	No change	No change	
	Steeply sloping hills with deep, fine soils	<i>Jhum</i> paddy (local land races)	No change	No change	
		Maize (local land races)	No change	No change	
	Moderately to gentle sloping hills	<i>Jhum</i> paddy (local land races)	No change	No change	

	slopes with deep loamy skeletal to fine loamy soils	Jhum-Maize (local land races)	No change	No change	
Delay by 6 weeks (3 rd May)	NA				
Delay by 8 weeks (1 st June)	NA				

2.1.1 .2 South west monsoon - normal (3rd week of May-Sept)

Conditions	Major Farming situation	Normal Crop / Cropping system	Change in crop / cropping system including variety	Suggested Contingency measures	
				Agronomic measures	Remarks on Implementation
Early season drought (delayed onset)	Moderately sloping on side slopes of hills with deep, fine loamy soils	<i>Kharif</i> maize (local land races)	No change		
		Terrace rice cultivation (local land races)	No change		
	Gently sloping, side slopes of hills with moderately shallow fine soils	<i>Kharif</i> maize	No change		
		Terrace rice cultivation (local land races)	No change	ICM	
	Steeply sloping, hills with deep fine soils	<i>Kharif</i> maize (local land races)	No change		
		Colocasia (local land races)	Summer vegetables	Mulching with local bio-mass (tree litter)	
	Moderately to gentle sloping hills slopes with deep loamy skeletal to fine loamy soils	<i>Kharif</i> maize (local land races)	No change		
		Terrace rice cultivation (local land races)	No change	ICM	

2.1.1 .3 South west monsoon - normal (3rd week of May-Sept)

Early season drought (delayed onset)	Major Farming situation	Normal Crop / Cropping system	Change in crop / cropping system including variety	Agronomic measures	Remarks on Implementation
Delay by 4 weeks (4 th week June)	Moderately sloping on side slopes of hills with deep, fine loamy soils	<i>Kharif</i> maize (local land races)	Local land races of maize Intercrop with Legumes (Soybean) and oilseeds (sesame) and local cucumbers	Mulching with local bio mass.	-
		Terrace rice cultivation (local land races)	Medium duration variety RCM-9, MTU-1010	ICM	
	Gently sloping , side slopes of hills with moderately shallow fine soils	<i>Kharif</i> maize (local land races)	Local land races of maize Intercrop with Legumes (Soybean) and oilseeds (sesame) and local cucumbers	Mulching with local bio- mass. throughout the cropping period	
		Terrace rice cultivation (local land races)	Medium duration variety Abishak	ICM	
	Steeply sloping, hills with deep fine soils	Terrace rice cultivation (local land races)	Medium duration variety Abishak	ICM	
		<i>Kharif</i> maize (local land races)	Local land races of maize Intercrop with Legumes (Soybean) and oilseeds (sesame) and local cucumbers	Mulching with local bio mass.	
	Moderately to gentle sloping hills slopes with deep loamy skeletal to fine loamy soils	<i>Kharif</i> maize (local land races)	Local land races of maize Intercrop with Legumes (Soybean) and oilseeds (sesame) and local cucumbers	Mulching with local bio mass.	
		Terrace rice cultivation (local land races)	Medium duration variety Abishak	ICM	
Delay by 6 weeks (1 st week July)	NA	NA			
Delay by 8 weeks (4 th week July)	NA	NA			

2.1.1.4 Monsoon- Normal

Condition			Suggested Contingency measures			
Early season drought (Normal onset)	Major Farming situation	Normal Crop/ cropping system	Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation	
Normal onset followed by 15-20 days dry spell after sowing leading to poor germination/crop stand etc.	Moderately sloping on side slopes of hills with deep, fine loamy soils	Kharif maize	I. If there is poor germination (Less than 30%) resowing II. Gap filling III. life saving irrigation if possible IV. Weeding	rain water harvesting as resource conservation technology, mulching with locally available bio mass, and earthing up		
		Ginger	I. If there is poor germination resowing of rhizomes II. inter-cultural operations.	In situ moisture conservation, mulching with locally available bio mass and life saving irrigation if possible		
	Gently sloping , side slopes of hills with moderately shallow fine soils	Jhum paddy	I. If there is poor germination (Less than 30%) re-sowing II. Keep Weed free	In situ moisture conservation, mulching with locally available bio mass	-	
	Steeply sloping, hills with deep fine soils	Terrace rice cultivation	No change		Transplanting of 30-35 Days old seedlings	-
		Maize	I. If there is poor germination (Less than 30%) re-sowing II. Gap filling III. Weeding	In situ moisture conservation, mulching with locally available bio mass	-	
		Ginger		Mulching		
	Moderately to gentle sloping hills slopes with deep loamy skeletal to fine loamy soils	Jhum paddy	I. If there is poor germination (Less than 30%) re-sowing II. Weeding	-	-	

2.1.1.5 Monsoon Normal

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures			
			Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation	
Mid season drought (Long dry spell consecutive 2 weeks rainless long dry)	Moderately sloping on side slopes of hills with deep, fine loamy soils	Kharif maize	Weeding/ intercultural operations etc.	rain water harvesting as resource conservation technology, mulching with locally available bio mass, and earthing up	Line dept. schemes/ RKVY -	
		Ginger	intercultural operations,weeding.	rain water harvesting as resource conservation technology, mulching with locally available bio mass, and earthing up		
	Gently sloping , side slopes of hills with moderately shallow fine soils	Terrace rice cultivation paddy	Foliar spray with 2 % urea and MOP	-		
		Ginger	Weeding/ intercultural operations etc.	rain water harvesting as resource conservation technology, mulching with locally available bio mass, and earthing up		
	Steeply sloping, hills with deep fine soils fine soils	Jhum paddy	Weeding Foliar spray with 2 % urea and MOP after rain	-		
		Maize	Weeding/ intercultural operations etc. Foliar spray with 2 % urea and MOP	rain water harvesting as resource conservation technology, mulching with locally available bio mass, and earthing up		
		Moderately to gentle sloping hills slopes with deep loamy skeletal to fine loamy soils	Jhum paddy	Weeding Foliar spray with 2 % urea and MOP		-
			Maize	Weeding/ intercultural operations etc. Foliar spray with 2 % urea and MOP		rain water harvesting as resource conservation technology, mulching with locally available bio mass, and earthing up

2.1.1.6 Monsoon Normal

Condition			Suggested Contingency measures		
Mid season drought (Long dry spell consecutive 2 weeks rainless long dry)	Major Farming situation	Normal Crop /cropping system	Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation
At flowering / fruiting stage	Moderately sloping on side slopes of hills with deep, fine loamy soils	Kharif, Maize,	Weeding/ intercultural operations etc. Life saving irrigation.	rain water harvesting as resource conservation technology, mulching with locally available bio mass, and earthing up	Line dept. schemes/ RKVY
		Ginger	life saving irrigation	rain water harvesting as resource conservation technology, mulching with locally available bio mass, and earthing up	
	Gently sloping , side slopes of hills with moderately shallow fine soils	Terrace rice cultivation paddy	Foliar spray with 2 % urea and MOP	-	
		Ginger	Weeding/ intercultural operations etc.	rain water harvesting as resource conservation technology, mulching with locally available bio mass, and earthing up	
	Steeply sloping, hills with deep fine soils fine soils	Jhum paddy	Weeding	-	
		Maize	Weeding/ intercultural operations etc.	rain water harvesting as resource conservation technology, mulching with locally available bio mass, and earthing up	
	Moderately to gentle sloping hills slopes with deep loamy skeletal to fine loamy soils	Jhum paddy	Weeding	-	
		Maize	Weeding/ intercultural operations etc.	rain water harvesting as resource conservation technology, mulching with locally available bio mass, and earthing up	

2.1.1.7 Terminal drought

Condition	Major Farming situation ^a	Normal Crop /cropping system	Suggested Contingency measures		
			Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation
Terminal drought (Early withdrawal of monsoon)	Moderately sloping on side slopes of hills with deep, fine loamy soils	Kharif, Maize,	Mulching Life saving irrigation if possible If grain filling is severely affected harvest for fodder	Land preparation for early rabi sowing of linseed, toria/pea	
		Ginger *	Mulching Harvest at physiological maturity	-	
	Gently sloping , side slopes of hills with moderately shallow fine soils	Terrace rice cultivation paddy	If grain filling is severely affected harvest for fodder	Land preparation for early rabi sowing of linseed, toria/pea	
		Ginger	Mulching Harvest at physiological maturity	-	
	Steeply sloping, hills with deep fine soils fine soils	Jhum paddy	If grain filling is severely affected harvest for fodder	Land preparation for early rabi sowing of linseed, toria/pea	
		Maize	Mulching and Life saving irrigation if possible Harvest at physiological maturity	-	
	Moderately to gentle sloping hills slopes with deep loamy skeletal to fine loamy soils	Jhum paddy	If grain filling is severely affected harvest for fodder	Land preparation for early rabi sowing of linseed, toria/pea	
		Maize	Mulching and Life saving irrigation if possible Harvest at physiological maturity	-	

2.2 Unusual rains (untimely, unseasonal etc) (for both rainfed and irrigated situations) NA

2.3 Floods: Not Applicable

2.4 Extreme events- Hailstorm

Extreme event type	Suggested contingency measure ^f			
	Seedling / nursery stage	Vegetative stage	Reproductive stage	At harvest
Hailstorm				
Tomato	NA	NA	NA	Harvest and value addition
Pineapple	NA	NA	NA	Harvest and value addition
Cucurbits	NA	Remove the affected plants and top dress with urea	NA	NA

* Other extreme events are not applicable in this district

Contingent strategies for Livestock, Poultry & Fisheries

2.5.1 Livestock

	Suggested contingency measures		
	Before the event	During the event	After the event
Drought/ Lean period (Oct-March)			
Feed and fodder availability	Encourage perennial fodder on bunds and waste land on community basis Establishing fodder banks, encouraging hedge row species for fodder crops	Utilizing fodder from perennial trees and Fodder bank reserves Transporting excess fodder from adjoining districts Use of non conventional fodders. Use of feed mixtures and feed blocks Culling unproductive livestock	Use of non conventional fodders. Use of feed mixtures and feed blocks Availing Insurance
Drinking water	Roof top water harvesting, Preserving water in the tank for drinking purpose.	Judicious use of water, Using preserved water in the tanks for drinking purpose, recycling of household used water. Chlorination of water.	Maintenance/cleaning of community reservoirs/ village ponds
Health and disease management	Insurance, Veterinary preparedness with medicines and vaccines, organizing vaccination camps and mineral supplementation	Conducting mass animal Health Camps and treating the affected one, mineral supplementation.	Culling sick animals and mineral supplementation
Floods	Not applicable		
Feed and fodder availability			
Drinking water			
Health and disease management			

Cyclone	Not applicable		
Feed and fodder availability			
Drinking water			
Health and disease management			
Heat wave and cold wave	Cold wave		
Shelter/environment management	Adoption of deep litter system for pig/poultry	Covering of open space with gunny bags, Warming of pen using heating bulb or any other source and Feeding of high energy feed	
Health and disease management	Deworming, hygiene and cleanliness of the floor of the pen	Apply appropriate medicine	

^s based on forewarning wherever available

2.5.2 Poultry

	Suggested contingency measures			Convergence/linkages with ongoing programs, if any
	Before the event	During the event	After the event	
Drought	-	-	-	-
Shortage of feed ingredients	Procurement and storage of feed ingredients, Establishing feed reserve Bank	Utilizing from feed reserve banks, nutritional supplementation to poultry	Nutritional supplementation to poultry	
Drinking water	Arrangement for drinking water, Roof top water harvesting , Preserving water in the tank for drinking purpose	Judicious use of water, providing B-complex and Vit.C in water	Supplementation of Vit. B-complex to be continued.	
Health and disease management	Insurance and Emergency Veterinary preparedness with medicines and vaccination to birds	Sanitation and Hygiene	Culling affected birds, Mass vaccination	
Floods	Not applicable			
Cyclone	Not applicable			
Heat wave and cold wave	Not applicable			

^a based on forewarning wherever available

2.5.3 Fisheries/ Aquaculture

	Suggested contingency measures		
	Before the event	During the event	After the event
1) Drought	-	-	--
A. Capture			
Marine	-	-	-
Inland	-	-	-
(i) Shallow water depth due to insufficient rains/inflow	-	-	-
(ii) Changes in water quality	-	-	-
(iii) Any other	-	-	-
B. Aquaculture			
(i) Shallow water in ponds due to insufficient rains/inflow	De-silting, repair of bunds of existing ponds, rain water harvesting, liming and adopt low stocking density, deepening of ponds by 1.5-2 meters, restrict use of Manures and fertilizers, Channelizing water to pond if possible, Maintain proper water quality	Integrated farming, air breathing fish to be practiced, avoid fertilization and manuring on supplementary basis, feeding should be minimum to avoid organic loading, short term aquaculture with medium and minor carps, Maintain proper water quality	Prepare pond for the next crop after early harvest, Maintain proper water quality
(ii) Impact of salt load build up in ponds / change in water quality	Rain water harvesting, deepening, de-silting of existing water bodies and removal of debris	Rain water harvesting, deepening, de-silting of existing water bodies and removal of debris	Control feeding to avoid waste accumulation and eutrophication
(iii) Any other			
2) Floods	Not Applicable	Not Applicable	Not Applicable
3. Cyclone / Tsunami	Not Applicable	Not Applicable	Not Applicable
4. Heat wave and cold wave	No change	No change	No change

^a based on forewarning wherever available