

State: Uttar Pradesh

Agriculture Contingency Plan for District: Mau

1.0 District Agriculture profile			
1.1	Agro-Climatic/Ecological Zone		
	Agro Ecological Sub Region (ICAR)	Eastern Plain, Hot Subhumid (moist) Eco-Region (13.1) & Northern Plain, Hot Subhumid (Dry) Eco-Region (9.2)	
	Agro-Climatic Zone (Planning Commission)	Middle Gangetic Plain Region (IV)	
	Agro Climatic Zone (NARP)	Eastern Plain Zone (UP-9)	
	List all the districts falling under the NARP Zone* (*>50% area falling in the zone)	Barabanki, Ambedkarnagar, Faizabad, Sultanpur, Azamgarh, Mau, Jaunpur, Varanasi, Gazipur, Ballia, Bhadohi	
	Geographic coordinates of district headquarters	Latitude	Longitude
		25°57' N	83°36' E
	Name and address of the concerned ZRS/ ZARS/ RARS/ RRS/ RRTTS	Directorate of Research, SAU, Kumarganj	
	Mention the KVK located in the district with address	KVK, Mau	

1.2	Rainfall	Normal RF(mm)	Normal Rainy days (number)	Normal Onset	Normal Cessation
	SW monsoon (June-Sep):	992.8	40	3 rd week of June	1 st week of October
	NE Monsoon (Oct-Dec):	61.2	2	-	-
	Winter (Jan- Feb)	33.6	5	-	-
	Summer (March-May)	29.7	3	-	-
	Annual	1117.3	50	-	-

1.3	Land use pattern of the district (latest statistics)	Geographical area	Cultivable area	Forest area	Land under non-agricultural use	Permanent pastures	Cultivable wasteland	Land under Misc. tree crops and groves	Barren and uncultivable land	Current fallows	Other fallows
	Area ('000 ha)	171.62	125.29	0.56	22.81	0.204	2.2	3.5	1.74	12.77	2.4

1.4	Major Soils (common names like red sandy loam deep soils (etc.,))*	Area ('000 ha)	Percent (%) of total Geographical area
	Loam/Clay Loam, Irrigated soils	70.13	62.62
	Rain fed Loam/clay loam soils	24.38	21.77
	Diara and river bed, Tauns & Ghaghra clay loam soils	3.67	3.28
	Deep (pond, reservoir and lake) clay soils	3.96	3.54
	Low land 30-45 cm depth	5.28	4.71
	Rain fed Sodic soils	1.28	1.14
	Irrigated Sodic soils	4.27	3.81

Source: districtprofile Uttar pradesh

1.5	Agricultural land use	Area ('000 ha)	Cropping intensity(%)
	Net sown area	125.29	172.4
	Area sown more than once	39.61	
	Gross cropped area	164.9	

1.6	Irrigation	Area ('000 ha)
	Net irrigated area	115
	Gross irrigated area	-

	Rainfed area	-		
	Sources of Irrigation	Number	Area ('000 ha)	Percentage of total irrigated area
	Canals	-	71.01	19.2
	Tanks	-	0.004	-
	Open wells	-	0.1	-
	Bore wells	17.5	174.05	47.25
	Lift irrigation schemes	-	-	-
	Micro-irrigation	-	-	-
	Other sources (please specify)	-	-	-
	Total Irrigated Area	-	368.3	-
	Pump sets	-	135	36.6
	No. of Tractors	-	-	-
	Groundwater availability and use* (Data source: State/Central Ground water Department /Board)	No. of blocks/ Tehsils	(%) area	Quality of water (specify the problem such as high levels of arsenic, fluoride, saline etc)
	Over exploited	-	-	-
	Critical	-	-	-
	Semi- critical	-	-	-
	Safe	-	-	-
	Wastewater availability and use	-	-	-
	Ground water quality			
*Over-exploited: groundwater utilization > 100%; critical: 90-100%; semi-critical: 70-90%; safe: <70%				

1.7 Area under major field crops & horticulture

1.7	Major field crops cultivated	Area ('000 ha)							
		Kharif			Rabi			Summer	Grand total
		Irrigated	Rainfed	Total	Irrigated	Rainfed	Total		
Rice	114.5	0.3	114.9	-	-	-	-	-	
Sorghum	-	0.9	0.9	-	-	-	-	-	
Maize	0.3	0.3	0.6	-	-	-	-	-	
Pigeon pea	-	3.3	3.3	-	-	-	-	-	
Wheat	-	-	-	118.061	0.066	118.127	-	-	
Chickpea	-	-	-	0.186	0.968	1.154	-	-	
Pea	-	-	-	4325	0.003	4.328	-	-	

	Horticulture crops - Fruits	Area ('000 ha)		
		Total	Irrigated	Rainfed
	-	-	-	-
	Horticulture crops - Vegetables	Total	Irrigated	Rainfed
	Potato	3998	3998	0
	Onion	0.1	0.15	0
	Others	6.9	6.6	0.2
	Medicinal and Aromatic crops	-	-	-
	Plantation crops	-	-	-
	-	-	-	-
	Eg., industrial pulpwood crops etc.	-	-	-
	Fodder crops	Total	Irrigated	Rainfed
	Jowar	0.94	0	0.9
	Bajra	0.03	0	0.03
	Maize	0.42	0.05	0.3
	Total fodder crop area	3.65	1.245	2.4

	Grazing land	-	-	-
	Sericulture etc	-	-	-
	Others (specify)	-	-	-

1.8	Livestock	Male ('000)	Female ('000)	Total ('000)	
	Indigenous	-	-	308.02	
	Non descriptive Cattle (local low yielding)	-	-	-	
	Improved crossbred cattle (Cow & Buffalo only)	-	-	214.08	
	Non descriptive Buffaloes (local low yielding)	-	-	-	
	Buffaloes	-	-	154.6	
	Goat	-	-	165.3	
	Sheep	-	-	598.2	
	Others (Camel, Pig, Yak, Horse, Monkey etc.)	-	-	10.2	
	Commercial dairy farms (Number)	-	-	-	
1.9	Poultry	No. of farms	Total No. of birds ('000)		
	Commercial	-	-		
	Backyard	-	-		
	Total		141.295		
1.10	Fisheries (Data source: Chief Planning Officer)				
	A. Capture				
	i) Marine (Data Source: Fisheries Department)	No. of fishermen	Boats		Storage facilities (Ice plants etc.)
			Mechanized	Non-mechanized	
			Mechanized (Trawl nets, Gill nets)	Non-mechanized (Shore Seines, Stake & trap nets)	
		-	-	-	-
	ii) Inland (Data Source: Fisheries Department)	No. Farmer owned ponds		No. of Reservoirs	No. of village tanks
		-		-	-
	B. Culture				

		Water Spread Area (ha)	Yield (t/ha)	Production ('000 tons)
	i) Brackish water (Data Source: MPEDA/ Fisheries Department)	-	-	-
	ii) Fresh water (Data Source: Fisheries Department)	-	-	-
	Others	-	-	-

1.11 Production and Productivity of major crops

1.11	Name of crop	Kharif		Rabi		Summer		Total		Crop residue as fodder ('000 tons)
		Production ('000 t)	Productivity (kg/ha)	Production ('000 t)	Productivity (kg/ha)	Production ('000 t)	Productivity (kg/ha)	Production ('000 t)	Productivity (kg/ha)	
Major Field crops (Crops to be identified based on total acreage)										
	Rice	292.8	2549	-	-	-	-	292.8	2549	-
	Sorghum	0.84	897	-	-	-	-	0.84	897	-
	Maize	0.8	1202	-	-	-	-	0.8	1202	-
	Pigeon pea	3.8	1132	-	-	-	-	3.8	1132	-
	Wheat	-	-	383.4	3246	-	-	383.4	3246	-
	Chick pea	-	-	1.19	1034	-	-	1.19	1034	-
	Pea	-	-	4.62	1068	-	-	4.62	1068	-
	-	-	-	-	-	-	-	-	-	-
Major Horticultural crops (Crops to be identified based on total acreage)										
Crop 1	-	-	-	-	-	-	-	-	-	-

1.12	Sowing window for 5 major field crops (start and end of normal sowing period)	Paddy	Maize	Pigeonpea	Wheat	Lentil
	Khariif- Rainfed	2 nd week of June-3 rd week of July	1 st week of June-4 th week of June	1 st week of July - 4 th week of July	-	-
	Khariif-Irrigated	4 th week of June-2 nd week of August	3 rd week of June-2 nd week of July	-	-	-
	Rabi- Rainfed	-	-	Early rabi- September -October	2 nd week of October - 2 nd week of November	1 st week of October – 3 rd week of October

	Rabi-Irrigated	-	-	-	2 nd week of November- 4 th week of December	-
--	----------------	---	---	---	-----------------------------------------------------------------------	---

1.13	What is the major contingency the district is prone to? (Tick mark)	Regular	Occasional	None
	Drought	-	√	-
	Flood	-	√	-
	Cyclone	-	-	√
	Hail storm	-	-	√
	Heat wave	-	√	-
	Cold wave	-	√	-
	Frost	-	-	√
	Sea water intrusion	-	-	√
	Pests and disease outbreak (specify)	-	√	-
	Others (specify)	-	-	-


1.14	Include Digital maps of the district for		
		Location map of district within State as Annexure I	Enclosed: Yes
		Mean annual rainfall as Annexure 2	Enclosed: Yes
		Soil map as Annexure 3	Enclosed: Yes

Annexure I




Agroclimatic Zones of U.P.

- | | |
|---|--------------------------|
| 1 | Dhabhar and Terai Zone |
| 2 | Western Plain Zone |
| 3 | Mid Western Plain zone |
| 4 | South Western Plain Zone |
| 5 | Central Plain Zone |
| 6 | Bundelkhand Zone |
| 7 | North Eastern Plain Zone |
| 8 | Eastern Plain Zone |
| 9 | Vidhya Zone |






Annexure II



Annexure III



2.0 Strategies for weather related contingencies

2.1 Drought

2.1.1 Rainfed situation

Condition			Suggested Contingency measures		
Early season drought (delayed onset)	Major Farming situation	Normal Crop / Cropping system	Change in crop / cropping system including variety	Agronomic measures	Remarks on Implementation
Delay by 2 weeks 1 st week of July	Deep loamy soils (tarai soils)	Rice	Rice Transplanting/Direct seeding of Medium and Short duration varieties of Rice Such as NDR-97, NDR-359,NDR-80,NDR-118, Baranideep etc.	Raise Staggered rice nursery should be grown at 15 days interval in small areas at least two times	
	Sandy loam soils	Maize	Maize	Intercropping/ mixed cropping of maize/sorghum/ Pearlmillet with long duration varieties of Pigeonpea	
		Pigeonpea	Pigeonpea	Sowing on raised beds, Intercropping with Maize/Blackgram/Greengram	

Condition		Suggested Contingency measures			
Early season drought (delayed onset)	Major Farming situation	Normal Crop / Cropping system	Change in crop / cropping system including variety	Agronomic measures	Remarks on Implementation
Delay by 4 weeks 3 rd week of July	Deep loamy soils (tarai soils)	Rice-Wheat	Rice-Wheat Transplanting/Direct seeding of Medium and Short duration varieties of Rice Such as NDR-97, NDR-359,NDR-80,NDR-118, Baranideep, Govind,Saket-4, Ratna,IR-36 and Pant-12 etc.	<ul style="list-style-type: none"> • Direct seedling of short duration varieties of Rice such as NDR-97, NDR-80, NDR-118, Saket-4 • Raise Staggered rice nursery should be grown at 15 days interval in small areas at least two times • Adopt SRI system of nursery raising • Transplanting of Rice (beyond 20th July) with 3-4 seedlings/hill to increasing the plant population of 60 hills/m², instead of 50 hills/m². • Pruning of over aged Rice seedlings for better establishment and optimum plant stand • Foliar spraying of 2.5 kg Urea + 2.5 kg Potash as to increase the drought tolerance in nursery / standing crops 	<ul style="list-style-type: none"> • Seed-drill under RKVY • Supply of seed through govt. agencies <i>ie.</i> NFSM,RKVY
	Sandy loam soils	Maize	Maize Maize-Prakash, Sartaj, Naveen, Tarun.	<p>Maintain weed free condition for soil moisture conservation</p> <p>Intercropping/ mixed cropping of maize with long duration varieties of Pigeonpea</p>	
		Pigeonpea	No change	<p>Sowing on raised beds</p> <p>Intercropping with Maize/Blackgram/Greengram</p>	

Condition	Major Farming situation	Normal Crop / Cropping system	Suggested Contingency measures		
			Change in crop / cropping system including variety	Agronomic measures	Remarks on Implementation
Delay by 6 weeks 1 st week of August	Deep loamy soils (tarai soils)	Rice-Wheat	Rice-Wheat Paddy: Short duration varieties of paddy such as NDR-97, NDR-80, NDR-118, Pant Dhan-12 should be transplanted/direct seeding.	Direct seeding of rice In case of late transplanting of rice (beyond 20 th July) planting should be dense by increasing the number of seedlings/hill from 2 to 3 to 3 to 4. Adopt SRI system of nursery raising Weeding and interculture Foliar spraying of 2.5 kg Urea + 2.5 kg Potash as to increase the drought tolerance in nursery / standing crops Life saving irrigation in transplanted rice	<ul style="list-style-type: none"> • Seed-drill under RKVY • Supply of seed through govt. agencies <i>ie.</i> NFSM, RKVY
	Sandy loam soils	Maize	Greengram/ Blackgram Greengram: T-44, Pant mung-1, Narendra mung-1 Blackgram : Narendra urd-1, Pant urd-25	Intercropping/ mixed cropping of Greengram/ Blackgram/ maize/sorghum/ Pearl millet with long duration varieties of pigeonpea Sorghum+green gram(2:2)	
		Pigeonpea	Pigeonpea: Bahar	Maize (Tipekhiya) in Pigeonpea(Narendra Arhar-1) crop in 1:1 row ratio Sowing on raised beds Intercropping with Maize/Blackgram/Greengram Pigeonpea+ Blackgram/Greengram (1:3)	

Condition			Suggested Contingency measures		
Early season drought (delayed onset)	Major Farming situation	Normal Crop / Cropping system	Change in crop / cropping system including variety	Agronomic measures	Remarks on Implementation
Delay by 8 weeks 3 rd week of August	Deep loamy soils (tarai soils)	Rice-Wheat	<p>Preference should be given for sowing of Pearl millet and Sesame</p> <p>Pearl millet: Pusa 322, 323(Hybrid) and WCC-75, Raj-171(Composite)</p> <p>Sesame: - Type-4, Type-78, Type-12</p> <p>Greengram : T-44, Pant mung-1, Pant mung-2, Samrat, Malviya, Janpriya, Malviya jyoti, Narendra mung-1</p>	<p>Direct sowing</p> <p>In case of late transplanting of rice(beyond 20th July) planting should be dense by increasing the number of seedlings/hill from 2 to 3 to 3 to 4.</p> <p>Foliar spraying of 2.5 kg Urea + 2.5 kg Potash as to increase the drought tolerance in nursery / standing crops</p> <p>Life saving irrigation in transplanted rice</p>	<ul style="list-style-type: none"> • Seed-drill under RKVY • Supply of seed through govt. agencies <i>ie.</i> NFSM,RKVY
	Sandy loam soils	Maize	<p>Blackgram : Narendra urd-1,Pant urd-25, Pant urd-19, Uttara, Type-9</p>	<p>Intercropping/ mixed cropping of Greengram/ Blackgram/maize/sorghum/ Pearl millet with long duration varieties of pigeonpea</p> <p>Land preparation for sowing of early rabi crops like potato,toria,lahi and mustard</p>	
		Pigeonpea	<p>September Pigeonpea Varieties Bahar, PDA-11, Pusa-9 should be done till 1st week of September.</p>	-	

Condition			Suggested Contingency measures		
Early season drought (Normal onset)	Major Farming situation	Normal Crop/cropping system	Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation
Normal onset followed by 15-20 days dry spell after sowing leading to poor germination/ crop stand etc.	Deep loamy soils (tarai soils)	Rice	<p>After seeding of rice if there is break of monsoon by 7 to 10 days and if seedling mortality is observed then re-sowing with the same variety</p> <p>Gap filling/transplanting in rice</p> <p>Using “Sanda” method, plant population can be maintained with sufficient number of tillers in late drought condition as to minimize the production losses</p>	<p>Weeding at critical stages</p> <p>Foliar praying of 2.5 kg Urea + 2.5 kg Potash as to increase the drought tolerance in nursery / standing crops</p> <p>Life saving irrigation</p> <p>Proper electricity monitoring/rostering system should be ensured in area for regular supply of electricity for pumping of water for life saving irrigation</p>	<ul style="list-style-type: none"> • Supply of inter cultural implements through RKVY • Farm ponds through IWSM programme • Pulse crop seeds supply through NFSM
	Sandy loam soils	Maize	<p>Ridge sowing</p> <p>Gap filling/ Thinning to maintain optimum plant population</p>	Leaf mulching to conserve the soil moisture	
		Pigeonpea	<p>Ridge sowing</p> <p>Gap filling/ Thinning to maintain optimum plant population</p>	Leaf mulching to conserve the soil moisture	

Condition			Suggested Contingency measures		
Mid season drought (long dry spell, consecutive 2 weeks rainless (>2.5 mm) period)	Major Farming situation	Normal Crop/cropping system	Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation
At vegetative stage	Deep loamy soils (tarai soils)	Rice	<p>Gap filling/transplanting in rice</p> <p>Foliar spraying of 2% urea to boost up the growth</p>	<p>Weeding as to conserve the residual soil moisture</p> <p>Leaf mulching to conserve the soil moisture</p> <p>Foliar spraying of 2.5 kg Urea + 2.5 kg Potash as to increase the drought tolerance in nursery / standing crops</p> <p>Life saving irrigation from the stored water during the rainy season.</p> <p>Proper electricity monitoring/rostering system should be ensured in area for regular supply of electricity for pumping of water for life saving irrigation</p>	
	Sandy loam soils	Maize	<p>Thinning to maintain proper distance between the plants.</p> <p>Frequent interculture</p> <p>Earthing up in Pigeonpea</p> <p>Foliar spraying of 2% urea to boost up the growth</p>	<p>Foliar spraying of 2% MOP to increase the resistance to drought</p> <p>Leaf mulching to conserve the soil moisture</p> <p>Conservation furrow</p> <p>Life saving irrigation</p>	
		Pigeonpea	-		

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation
At flowering/ fruiting stage	Deep clay loam soils	Rice	Foliar spraying of 2% urea to boost up the growth	<p>Weeding as to conserve the residual soil moisture</p> <p>Leaf mulching to conserve the soil moisture</p> <p>Foliar spraying of 2.5 kg Urea + 2.5 kg Potash as to increase the drought tolerance in nursery / standing crops</p> <p>Life saving irrigation from the stored water during the rainy season.</p> <p>Proper electricity monitoring/rostering system should be ensured in area for regular supply of electricity for pumping of water for life saving irrigation</p>	
		Maize	Thinning to maintain proper distance between the plants.	Foliar spraying of 2% MOP to increase the resistance to drought	
		Pigeonpea	<p>Frequent interculture</p> <p>Earthing up in Pigeonpea</p> <p>Foliar spraying of 2% urea to boost up the growth</p>	<p>Leaf mulching to conserve the soil moisture</p> <p>Conservation furrow</p> <p>Life saving irrigation</p>	

Condition			Suggested Contingency measures		
Terminal drought (Early withdrawal of monsoon)	Major Farming situation	Normal Crop/cropping system	Crop management	Rabi Crop planning	Remarks on Implementation
	Deep clay loam soils	Rice	<p>Alternate management of irrigation should be ensured for provide life saving irrigation</p> <p>Proper electricity monitoring/ rostering system should be ensured in area for regular supply of electricity for pumping of water for life saving irrigation</p>	<p>Better pulverization should be made for conservation of soil moisture following by planking for sowing of early rabi crops like toria and potato etc..</p> <p>Toria variety- type-9, type-36, PT-303, PT-30 and ageti Rai should be sown in 1st week of September while Bhawani variety can be sown in 2nd week of September.</p> <p>In fallow fields to sow Ageti rai, potato varieties like Kufri Ashoka, Kufri Chandra mukhi and other vegetable crops like spinach,reddish coriander etc.</p>	
		Maize	<p>Harvesting of intercrop at physiological maturity (Maize, Blackgram and Greengram)</p> <p>Earthing up of Pigeonpea</p> <p>Harvesting of green cobs (maize) and sell in market and remaining portion will be used for fodder.</p> <p>Life saving irrigation to pigeonpea if possible.</p>	<p>Better pulverization should be made for conservation of soil moisture following by planking for sowing of early rabi crops like toria and potato etc..</p> <p>Toria variety- type-9, type-36, PT-303, PT-30 and ageti Rai should be sown in 1st week of September while Bhawani variety can be sown in 2nd week of September.</p>	
		Pigeonpea		-	

2.1.2 Drought - Irrigated situation

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delayed release of water in canals due to low rainfall	Deep clay loam soils	Rice – Wheat / Pea/ Lentil	Short duration rice varieties- NDR 97, Ratna, Narendra 118, Narendra 97, Pant Dhan 12, HUR 105, Induri Sambha, HUR 2-1, HUR-3022 to be grown under aerobic condition.	Community nursery Direct seeding in small beds. Use of micro-irrigation systems viz. sprinkler & sub-surface irrigation.	Breeder's seed will be supplied by BHU and NDUAT, Faizabad. Seed drills RKVY and supply of seeds NFSM
Limited release of water in canals due to low rainfall	Sandy clay loam soils	Rice – Wheat / Pea/ Lentil	Rice\ Maize \ Sorghum Grow short duration aerobic rice such as NDR 97, NDR 118, Govind, Vandana, Varanideep, Susk Samrat , HUR 105 Maize: Malviya hybrid Makka-2, Naveen & Jaunpuri Pearl millet : WCC 75, Raj 171, Pusa 23 Sorghum: CSH-16, CHS-9, CHS-14, CSV-13 & CSV-15 should be grown on ridges for fodder/grain purposes.	Community nursery, Direct seeding in small beds. Use of micro-irrigation systems viz. sprinkler & sub-surface irrigation.	
Non release of water in canals under delayed onset of monsoon in catchment	Sandy clay loam soils	Rice – Wheat / Pea/ Lentil	Shift to only aerobic rice Or Rice may be replaced by pulses Greengram: Pant Mung -8, PDM-11, Samrat, Jyoti, Jagriti, Janpriya, Jan Chetana & Jan Kalyani Blackgram: Type 9, Pant U 19, Pant U 35, Narendra Urd 1 & Azad Urd-3 Sesame :Type 4, T-12, T-13, Shekhar, GT1, TC 25 & TC 289	Direct seeding in small beds. Use of micro-irrigation systems viz. sprinkler & sub-surface irrigation.	

Lack of inflows into tanks due to insufficient /delayed onset of monsoon	Sandy clay loam soils	Rice – Wheat / Pea/ Lentil	Sorghum\ Pearl millet	Conservation tillage, Sowing of Pearl millet & Sorghum for grain purposes at 45 cm on ridges. Foliar application of 2% MOP Use of mulches (straw/dust).	
Insufficient groundwater recharge due to low rainfall	Sandy clay loam soils	Rice – Wheat / Pea/ Lentil	Rice should be replaced with pulses (green gram & black gram), oilseeds (Sesame) in <i>Kharif</i> and wheat by Chickpea & lentil in <i>Rabi</i> season.	Direct seeding in small beds.	

2.2 Unusual rains (untimely, unseasonal etc) (for both rainfed and irrigated situations)

Condition	Suggested contingency measure			
	Vegetative stage	Flowering stage	Crop maturity stage	Post harvest
Continuous high rainfall in a short span leading to water logging	-	-	-	-
Rice	Provide drainage	Proper bunding, drain out excess water	Harvesting at physiological maturity	Shift to safer place
Wheat	Provide drainage	Drain out excess water	Harvesting at physiological maturity	Shift to safer place
Pigeonpea	Provide drainage and Practice of sowing on ridges	Make inter-row furrow to Drain out excess water	Harvesting at physiological maturity	Shift to safer place
Heavy rainfall with high speed winds in a short span²	-	-	-	-
Outbreak of pests and diseases due to unseasonal rains				
Rice, Wheat, Chickpea, Pigeonpea, Pearl millet	Need based plant protection (integrated pest and disease management)	Need based plant protection (integrated pest and disease management)	Need based plant protection (integrated pest and disease management)	Safe storage against stored grain pest and diseases

2.3 Floods

Condition	Suggested contingency measure			
	Seedling / nursery stage	Vegetative stage	Reproductive stage	At harvest
Transient water logging/ partial inundation¹				
Rice	<ul style="list-style-type: none"> • Arrangement of Drainage channel • Drainage of water from the rice fields 	<ul style="list-style-type: none"> • Removal of excess water 	<ul style="list-style-type: none"> • Foliar spray of 5% urea 	-
Maize	<ul style="list-style-type: none"> • Drainage of water • Creation of surface drains at appropriate places to avoid water logging 	-	-	-
Horticulture				
Continuous submergence for more than 2 days²	-	-	-	-
Rice	<ul style="list-style-type: none"> • Drainage of excess water through drainage channel • Transplanting of deep water rice –Madhupur, Jalmagn, Jalpriya, Jalnidhi, Awarodhi 	<ul style="list-style-type: none"> • Just after finishing of floods, topdressing of urea could be ensured in the field 	<ul style="list-style-type: none"> • Foliar spray of 5% urea 	<ul style="list-style-type: none"> • Preference should be given for planting of Autumn Sugarcane in the month of Oct so that their grand growth completed to the maxi. Extent prior to floods. • Planting of Sugarcane on raised beds instead of flat bed. • Emphasis could be given for cultivation of Toria, Urd, Mung/Sunflower
Horticulture				
Sea water intrusion	Not applicable			

2.4 Extreme events: Heat wave / Cold wave/Frost/ Hailstorm /Cyclone

Extreme event type	Suggested contingency measure ^f			
	Seedling / nursery stage	Vegetative stage	Reproductive stage	At harvest
Heat Wave				
Rice	• Frequently supply of fresh water	-	-	-
	• Avoid irrigation during hot period	• Micro-irrigation	-	-
Cold wave	•	•		
Wheat	Provide light irrigation	Provide light irrigation	Provide light irrigation	-
Pigeonpea	Mulching	Light irrigation for survival	Light irrigation for survival	Harvesting at physiological maturity
Frost	Not applicable			
Hailstorm	Not applicable			
All the crops	Re sowing	Re sowing of Catch crop	Harvest for fodder	Pre Harvesting
Cyclone	Not applicable			

2.5 Contingent strategies for Livestock, Poultry & Fisheries

2.5.1 Livestock

	Suggested contingency measures		
	Before the event ^g	During the event	After the event
Drought			
Feed and fodder availability	• Storage of straw and silage in Silo pit according to population of animal	-	-
Drinking water	• Maintenance and inspection of Tubewells, Hand pumps, Ponds, Tanks etc.	-	-
Health and disease management	• Vaccination of animals against FMD, HS, B.Q. and Dewarming	-	-
Floods			
Feed and fodder availability	• Increase the area of fodder	• Availability of safe place	• Sowing of rabi fodder crops

		for the animals	- Berseems, Lucerne, Oat and other rabi crops
Drinking water	•Crops according to population and their storage	•Distribution of stored feed and fodders according to the population of affected	• Drain of infected stored water and supply of fresh water for drinking.
Health and disease management	•Arrangement of clean drinking water in sufficient water in growth	•Provide neat & clean drinking water	• Proper treatment of affected (animals vaccination & Dewarming)
Cyclone	Not applicable		
Feed and fodder availability	•Arrangement of clean drinking water in sufficient water in growth	•Organize health camp regularly	-
Drinking water	-	-	-
Health and disease management	-	-	-
Heat wave and cold wave			
Shelter/environment management	•Shelter house/Farm house should not face directly	•Proper availability of shelter, drinking water and feeds & fodder as per need of the animals	•Provide shelterbelts of good quality materials
Health and disease management	•Ensure the availability of drinking water and as well as electrolytes	-	•Routine health check up by veterinary doctors

^s based on forewarning wherever available

2.5.2 Poultry

	Suggested contingency measures			Convergence/linkages with ongoing programs, if any
	Before the event	During the event	After the event	
Drought				
Shortage of feed ingredients	-	-	-	-
Drinking water	• Deep tubewell provide clean drinking	•Provide the drinking water	•Provide the drinking water	-
Health and disease management	• Vaccination against infectious diseases	•Vaccination	•Vaccination for infectious diseases such as- Ranikhet, infectious	-

			Coryza, IBD, ILT	
Floods				
Shortage of feed ingredients	<ul style="list-style-type: none"> • Inspection of established Tubewell & other water sources 	<ul style="list-style-type: none"> • Provide the drinking water 	<ul style="list-style-type: none"> • Provide the drinking water 	-
Drinking water	<ul style="list-style-type: none"> • Vaccination against infectious diseases 	<ul style="list-style-type: none"> • Vaccination 	<ul style="list-style-type: none"> • Vaccination for infectious diseases such as- Ranikhet, infectious Coryza, IBD, ILT 	-
Health and disease management	-	-	-	-
Cyclone	Not applicable			
Shortage of feed ingredients	-	-	-	-
Drinking water	-	-	-	-
Health and disease management	-	-	-	-
Heat wave and cold wave				
Shelter/environment management	<ul style="list-style-type: none"> • Arrangement of proper shelter and cooler/heater to maintain the proper temp. of the shelter house 	<ul style="list-style-type: none"> • Maintenance of surrounds temp. and prevent the birds from direct exposure of heat/ cold waves 	<ul style="list-style-type: none"> • Heat check up 	-
Health and disease management	<ul style="list-style-type: none"> • Vaccination 	<ul style="list-style-type: none"> • Vaccination 	<ul style="list-style-type: none"> • Vaccination • Availability of neat & clean water 	-

^a based on forewarning wherever available

2.5.3 Fisheries/ Aquaculture

	Suggested contingency measures		
	Before the event	During the event	After the event
1) Drought	-	-	-
A. Capture	-	-	-

Marine	-	-	-
Inland	Arrange for alternative water resources	Sell the produce at minimum acceptable size to the consumer	Lime Application
(i) Shallow water depth due to insufficient rains/inflow	Stocking of Air breathing	-	-
(ii) Changes in water quality	-	Increased water temperature	-
(iii) Any other	-	Decrease dissolve oxygen	-
B. Aquaculture	-	-	-
(i) Shallow water in ponds due to insufficient rains/inflow	Arrange for alternative water resources	Minimum disturbance to the fish i.e. minimum fishing activities	Maintain the pond properly by liming, manuring and fertilization
(ii) Impact of salt load build up in ponds / change in water quality	-	-	-
(iii) Any other	-	-	-
2) Floods	-	-	-
A. Capture	-	-	-
Marine	-	-	-
Inland	Harvest the large size fish	Protect the escape of fish	Manage the inlet, outlet structures along with pond land
(i) No. of boats / nets/damaged	-	-	-
(ii) No.of houses damaged	-	-	-
(iii) Loss of stock	-	-	-
(iv) Changes in water quality	-	-	-
(v) Health and diseases	-	-	-
B. Aquaculture	-	-	-
(i) Inundation with flood water	Make 2.5 m high bylonnet bundry on the band of pond	Check for outlet to remain open	Close outlet and open inlet
(ii) Water contamination and changes in water quality	-	Close inlet and divert water receiving channel	Treatment of water with Alum and $KmnO_4$
(iii) Health and diseases	-	-	Feeding, liming, manuring and

			fertilization of ponds
(iv) Loss of stock and inputs (feed, chemicals etc)	-	-	-
(v) Infrastructure damage (pumps, aerators, huts etc)	-	-	-
(vi) Any other	-	-	-
3. Cyclone / Tsunami	Not applicable		
4. Heat wave and cold wave			
A. Capture	-	-	-
Marine	-	-	-
Inland	-	-	-
B. Aquaculture	-	-	-
(i) Changes in pond environment (water quality)	-	-	-
(ii) Health and Disease management	-	-	-
(iii) Any other	-	-	-

^a based on forewarning wherever available